INFLUENCE OF MOTHER BULB NITROGEN ON SUBSEQUENT NITROGEN UPTAKE IN TULIPS

M. Amano
 Vegetable and Ornamental Crops Research Station
 Ano, Mie,
 Japan

STABY

Abstract

Influence of the nitrogen concentration of the mother bulb (planting bulb) was investigated by the experiment using 15N tracer technique. Mother bulb nitrogen had no significant effect either on the growth of above-ground parts at flowering or on the bulb yield at lifting. But both nitrate and ammonia nitrogen uptakes significantly decreased with increasing nitrogen concentration of the mother bulb.

Distribution factor of mother bulb nitrogen to each part of plant changed only slightly as nitrogen concentration of mother bulb increased. Regardless of mother bulb nitrogen, about 15 % of nitrogen originated from the mother bulb was transferred into the flower and removed by defloration, and finally 70 % was redistributed into daughter bulb and harvested.

From the above, it was clarified that there might be the inverse correlation between the nitrogen concentration of the mother bulb and subsequent uptake of nitrogen, and high recovery percentage could be obtained for mother bulb nitrogen in tulip.

1. Introduction

Nitrogen fertilization for tulip bulb production has a great influence upon not only the growth and bulb yield during the season's growth but also nitrogen content of daughter bulb used for planting in succesive year (Tsutsui et al.,1967,Amano et al., 1981). It is also important that nitrogen level of bulb is closely related to forcing ability (Cheal et al., 1964, Kabe, 1980, Munk, 1980) and infection of deseases (Tsutsui et al., 1967). Furthermore insecurity of nitrogen application effects is caused by remarkable leaching by nitrification in the soil (Hagiya et al., 1966, Tsutsui et al., 1969, Baba, et al., 1983).

There are few reports which give effects of planting-bulb nitrogen on bulb yield in successive year. The previous investigation showed that there were some cases where there might be the inverse correlation between nitrogen levels in mother bulbs and bulb yields (Amano et al., 1981).

The investigation reported here was designed to clarify the effects of mother bulb nitrogen on the subsequent nitrogen uptake and the distribution of various nitrogens, originated from mother bulb, absorbed from nitrate nitrogen and from ammonia nitrogen, by the experiment using $15_{\rm N}$ tracer technique.

2. Material and methods

Seed bulbs for the experiment were lifted from five different stocks of tulip Rose Beauty, to which the top dressing of nitrogen with 0, 4,

Acta Horticulturae 177, 1985 Flower Bulbs IV 8, 12, and 16 Kg / 10a were applied in the spring 1979. For each stock, planting mother bulbs of 9 cm in circumstance (9 cm grade) were chosen with bulb weight ranging from 15.5 to 15.8 g. Nitrogen concentration of planting mother bulb selected from each stock was 0.99, 1.22, 1.36, 1.64, and 1.87 % respectively.

Three mother bulbs from each stock were planted into a 1/2000 Wagner pot with sand. These plots comprised five levels of mother bulb nitrogen and each plot was consisted of ten pots.

Sand culture was made in an unheated plastic film house, with a supply of nutrient solution during the periods from 11 March to 31 May. Composition of the nutrient solution used in sand culture was given in

Table 1. Sodium nitrate included 4.64 atom % excess $15_{\rm N}$ as a nitrate form and ammonium sulfate included 2.19 atom % excess $15_{\rm N}$ as an ammonia form were applied. The solution containing both nitrogen form in the ratio of 1 : 1 was used, and the concentration of total nitrogen element was 100 ppm.

Samples of 12 plants at the flowering time (9 May) and daughter bulbs 18 plants at the lifting time (15 June) were taken for nitrogen analysis(Amano et al., 1984, 1985).

3. Results

3.1. Effects on the growth and the bulb yield

The growth of above-ground parts at flowering only slightly decreased with increasing the nitrogen concentration of mother bulb (Table 2). There was no clear effect on total bulb yield.

3.2. Effects on nitrogen content in each part of plant at flowering

There were downward trends in nitrogen contents of flowers, 1st leaves and roots and on the contrary upward trends in those of daughter bulbs and mother bulbs, with increasing mother bulb nitrogen. Mother bulb nitrogen resulted in having no clear effect on total nitrogen contents of whole plant at flowering (Table 3).

3.3. Effects of subsequent nitrogen uptake

The partition of nitrogen content between various nitrogens; originated from mother bulb, absorbed from $\mathrm{NH}_4-\mathrm{N}$ and from $\mathrm{NO}_3-\mathrm{N}$, were indicated in Figure 3. The percentage of nitrogen from mother bulb consistently increased in all parts of plant with increasing mother bulb nitrogen and there were outstanding decrease in nitrogen from both $\mathrm{NH}_4-\mathrm{N}$, which was more influenced by mother bulb nitrogen.

With regard to the whole plant at flowering, the nitrogen uptake from NO₃-N significantly decreased with increasing mother bulb nitrogen , as from NH₄-N (Table 4 , Figure 1). With regard to the daughter bulb at lifting, the nitrogen content from NO₃-N significantly deceased with increasing mother bulb nitrogen, but there was only a small effect on that from NH₄-N (Table 4, Figure 2).

3.4. Recovery of mother bulb nitrogen

Distribution factor of mother bulb nitrogen to each part of plant changed only slightly as nitrogen concentration of mother bulb increased. Each mean percentage of distribution to flower, leaves, stem, daughter bulbs, roots or mother bulb was 15, 36, 10, 26, 8, or 6 respectively (table 5). Recovery rate of mother bulb nitrogen into lifted bulbs was around 70 % regardless of mother bulb nitrogen.

4. Discussion

It was consistent with the results reported by Nishii (1963), Cheal (1963), Tsutsui (1967, 1969) and many other workers that there were clear effects of nitrogen supply on bulb weight and bulb nitrogen content in propotion to nitrogen level in that year.

However, various results were seen on those effects in the following year, despite the generally considered view that tulips have aftereffect of the nitrogen fertilization in the previous generation. Amaki (1960) found that in the large seed bulb, the yield of bulb was varied depending on the amount of nitrogen fertilizer supplied to the previous generation, but in the small seed bulb, such an aftereffect was not recognize. Tsutsui (1967) reported that the nitrogen supply at later stage had no aftereffect and Amanc (1981) reported that there were the inverse correlation between nitrogen levels in mother bulb and bulb yields. It was reported by Sekher (1981) that the application of nitrogen in the spring significantly increased the nitrogen content of bulbs, but, once again the application of nitrogen in the autumn had no effect. The results in this investigation showed that mother bulb nitrogen had no clear effect on the growth and bulb yield, nevertheless both nitrate and ammonia nitrogen uptakes significantly decreased with increasing nitrogen concentration of the mother bulb.

It is likely that such disagreement for aftereffect of nitrogen application as in the previous papers is caused by the inverse correlation between mother bulb nitrogen and subsequent nitrogen uptake. Furthermore, aftereffect might be also influenced by the later absorption as described by Tsutsui (1967).

It was proved that distribution factor of nitrogen originated from mother bulb to each part of plant changed only very slightly as nitrogen level of mother bulb increased, and 15 % of those was transferred into the flower which removed by defloration and finally 70 % was redistributed into daughter bulbs.

Large quantities of nitrogen originated from mother bulb would be subsequently redistributed into daughter bulbs at the lifting time, as the similar studies by Irobe (1966) and Schmalfeld (1965).

References

Amaki, W. and Hagiya, K., 1960. Studies on fertilizer supply to tulips.
II. The differences in growing of tulips bulbs produced by supplying
different amounts of fertilizers in preceding generation, during the forcing and field culture. J. Jap. Soc. Hort. Sci. 29:239-246. (In Japanese
with English summary).

Amano, M. and Nezu, M., 1981. Studies on the nitrogen fertilization of tulips. I. Effects of seed bulb nitrogen on the bulb yield. Abstract of Spring Meeting, Jap. Soc. Hort. Sci.(1981):525. (In Japanese).

Amano, M. and Imai, F., 1984. Studies on the nitrogen fertilization of tulips. I. Effects of mother bulb nitrogen on the subsequent uptake of 15NO₃-N and ¹⁵NH₄-N. Ibid.(1984):304-305. (In Japanese).

Amano, M. and Imai, F., 1985. Studies on the nitrogen fertilization of tulips. III. The traces of nitrogen originated from mother bulb using 15N tracer technique. Ibid.(1985):382-383. (In Japanese).

Baba, A., Ikarashi, T. and Yokoyama, S., 1983. Simple lysimeter test of

A 1.4.4. A.A.

the leaching of fertilizers in tulip cultivation. 2. Effects of controlling nitrification in the loam soil. Bull. Facul. Agric. Niigata Univ. 35:19-36. (In Japanese with English summary).

Cheal, W. F. and Hewitt, E. J., 1963. Effects of mineral nutrition on the production of tulip bulbs. Ann. appl. Biol. 52:493-592.

- Cneal, W. F. and Hewitt, E. J., 1964. The effects of rates of supply of nitrogen, phosphorus, potassium and magnesium on leaf and stem growth, flowering and topple of Golden Harvest and Elms tulip. Ann. appl. Biol. 53:477-484.
- Hagiya, K. and Amaki, W., 1966. Nutritional studies on tulips. N. The leaching of three major elements from the soil during the growing season. J. Jap. Soc. Hort. Sci. 35:107-114. (In Japanese with English summary).
- Irbe, A., Obata, S. and Yanagisawa, M., 1966. Studies on the nutrition of tulips (1). Transport of mother bulb nitrogen. Bull. Toyama Agric. Exp. Stn. 1:147-153. (In Japanese).
- Kabe, H., 1980. The influence of concentration of mineral nutrients on the forcing-ability of the tulip bulbs. Bull. Niigata Hortic. Exp. Stn. 10:15-29. (In Japanese with English summary).
- Munk, W. J. de, Hoogeterp, P. and Slootweg, G., 1980. Effects of nitrogen dressing on flower-bud blasting in tulips during forcing. Acta Horticulturae 109:81-87.
- Nishii, K. and Tsutsui, K., 1963. Studies on the nitrogen nutrition of tulips. I. Effects of the time of nitrogen supply on the absorption of nutrients and on growth and yields. J. Jap. Soc. Hort. Sci. 32:65-73. (In Japanese with English summary).

Schmalfeld, H. W. and Carolus, R. L., 1965. Nutrient redistribution in the tulip. Proc. Am. Soc. Hort. Sci. 86:701-707.

- Sekhar, N. N. C.; 1981. Nitrogen for tulip bulb production. Scientific Hort. 32:104-109.
- Tsutsui, K., Nishii, K. and Toyoda, T., 1967. Studies on the nitrogen nutrition of tulips. III. Effects of nitrogen top-dressing insucceeding crop system after paddy. Bull. Toyama Agric. Exp. Stn. 2: 129 -136. (In Japanese).
- Tsutsui, K. and Toyoda, T., 1969. Studies on the nitrogen nutrition of tulips. N. Effects of the top-dressing of nitrogen in the planting year. Tbid. 3:19-25. (In Japanese with English summary).

Table 1 -	Composition	of the	nutrient	solution	used	in
	sand culture	2				

ن جي هي هه هه ها ها ها ها ها خل خو هه جي خو	و این برد برد می دو دو می موجود این می دو	
Element	Conc. of element	Chemicals used
N	100 ppm	$\operatorname{Na^{15}NC}_{3} + (\operatorname{NH}_{4})_{2} \operatorname{SO}_{4}$
		$\operatorname{NaNO}_3 + ({}^{15}\operatorname{NH}_4)_2 \operatorname{SO}_4$
P	30	NaH2POL.2H20
К	100	K ₂ SO _L
Ca	80	CaCl ₂ ·2H ₂ O
Mg	50	MgSOL·TE2O
Mn	0.5	MnSO ₁ ·4E ₂ O
В	0.5	H ₃ BO ₃
Zn	0.1	ZnSO _L ·TE ₂ O
Cu	0.01	CuSO _L ·5E ₂ O
Мо	0.01	(NHL) 6 Mo 024 · 4H20
Fe	0.55	Fe-EDTA

pH : 5.8-6.0

:

Table 2 - Effect of nitrogen concentration of mother bulb on the growth at flowering (9 May) and bulb yield at lifting in tulip, Rose Beauty (9 cm grade planting bulb)

N % of mother bulb	Growth o	of tops (c	m)	Bulb yi	Bulb yield (g/plant)			
	Stem length	lst leaf length	lst leaf width	Main bulb	Lateral bulb	Total		
0.99	30.9	13.1	10.0	26.0	6.9	32 . 9		
1.22	29.5	13.6	10.2	25.9	7.3	33.2		
1.36	28.7	13.0	9.8	22.9	8.8	31.8		
1.64	28.4	13.2	9.6	24.4	9.0	33.4		
1.87	24.1	12.8	9.0	22.1	9.2	31.3		

427

and a second second

Table 3 - Effects of nitrogen concentration of mother bulb on nitrogen content (mg/plant) in the parts of plant of tulip , Rose Beauty at flowering (9 cm grade planting bulb)

Ç.,

N % of mother	Flower	Leaf				Stem	Daughter	Root	Mother	Total
bulb		lst	2nd	3rd	Total		DUTD		bulb .	
0.99	33	28	19	16	63	21	 41	 14	6	179
1.22	31	27	18	18	63	20	43	13	7	177
1.36	32	26	19	19	64	21	48	13	7	185
1.64	30	26	19	20	65	20	54	10	9	188
1.84 	27 	24	16 	17	57	18	54	11	9	176

Table 4 - Effects of nitrogen concentration of mother bulb on the contents of various nitrogen ; originated from mother bulb, absorbed from NO3-N and NH4-N, in tulip, Rose Beauty

N % of mother bulb	N mg / at flow	whole g vering	olant		N mg of at lift	daughter bulbs/plant			
	Mother bulb N	N0 ₃ -N	NH ₁ -N	Total	Mother Bulb N	NO ₃ -N	NH4-N	Total	
0.99	65	67	 46	 178	 53	 49			
1.22	80	56	41	177	75 71	.ጋ ኴ7	30	150	
1.36	112	53	20	185	84	36	בב ג)י	152	
1.64	128	46	13	138	104	36	30	170	
1.87	124	36	14	176	100	31	36	167	

• • •

and any high and an and a second second

Table 5 - Partition of mother bulb nitrogen between the parts of plant at flowering and recovery of mother bulb nitrogen at lifting in tulip, Rose Beauty

÷

N % of mother bulb	% of m	other]	bulb N a	% of N					
	Flower	Leaf	Stem	Daughter bulb	Root	Mother bulb	Whole plant	into lifted	• • •
0.99	1ć	37	10	20	10	.7	100	74	
1.22	<u>1</u> L .	38	9	23	9	7	100	72	
1.36	15	36	10	25	8	ć	100	75	
1.64	1ó	34	10	29	5	6	100	71	
1.87	15	32	10	31	6	б	100	56	
Mean	15	36	10	26	8	6	100	70	











Figure 3 - Influence of nitrogen concentration of mother bulb on the partition of nitrogen content between various nitrogen () originated from mother bulb,) absorbed from NH4-N and 1000 from NO3-N) in the parts of plant of tulip, Rose Beauty at flowering