

Potential Salad Crops for Cultivation

Usha Rani Palaniswamy

Graduate Student, Department of Plant Science

Reprinted from the Connecticut Flower Growers Newsletter

A number of epidemiological and clinical studies have demonstrated the benefits of vegetables and fruits in the human diet in the prevention of several major diseases including cancer and heart disease. Vegetables and fruits contain chemicals that prevent the occurrence and progression of these major diseases. The National Cancer Institute and the American Heart Association recommend that we Strive for Five—five servings of fruits and vegetables a day for a healthier

of our food needs. Perhaps it's time to broaden our food base and look for alternative and better sources of nutraceuticals.

Watercress is a leafy perennial that grows well at temperatures of 20° to 25°C (68° to 77°F). It is a salad crop commonly used as a pot herb. It is easily propagated by seed, stem or terminal shoot cuttings and requires about 25 to 30 days from seedling transplant to harvest. It contains a chemical, phenethyl isothiocyanate (PEITC), that can prevent cancers caused by carcinogenic nicotine compounds. Watercress is also a rich source of vitamins A and C.

"Vegetables and fruits contain chemicals that prevent the occurrence and progression of major diseases including cancer."

life. The nutritional importance of vegetables has long been recognized. Currently, there is an increased awareness among the general public of the health benefits of the specific chemicals in vegetables and fruits that are sometimes referred to as "nutraceuticals". Among the 13,000 known edible plants, less than 20 are currently being used to provide most

Purslane is an annual succulent that grows well at temperatures of 22° to 27°C (71° to 81°F) and has a high growth rate and water use efficiency. It is a relative of the common spring annual flowering *Portulaca* and a common prostrate weed in the fields. There are two upright types, a green-leafed type and a golden leafed type that are cultivated as salad greens. All types are propagated easily by seeds, stem or shoot cuttings and take about 20 to 25 days from seedling transplant to harvest. Purslane is an excellent source of the essential omega-3 fatty acid known as alpha-linolenic acid. The role of this

omega-3 fatty acid in normal human growth, development and disease prevention has been well established. Research indicates that omega-3 fatty acids will eventually receive the attention and broad recommendations now given to food fiber.

Although watercress is a common salad crop, its cultivation has declined over time, and purslane is still considered a weed with a nutritive potential. However, it is encouraging to note that purslane is being considered for cultivation by the USDA and is regarded as the Power Food of the Future because of its high omega-3 fatty acid and vitamin E contents.

I am conducting research in the Department of Plant Science, University of Connecticut, under the direction of Dr. R. McAvoy, to identify and characterize the environmental factors (light intensity, day length, temperature, mineral nutrition, etc.) that would optimize the nutraceuticals in the edible parts of purslane and watercress.

To date, we have found that leaves of Watercress plants grown in a hydroponic medium containing 200 ppm nitrogen and 128 ppm sulfur contain 84% more PEITC than those grown in a medium containing 200 ppm nitrogen and 64 ppm sulfur. The leaves of plants grown in a medium containing 200 ppm nitrogen and 192 ppm sulfur contain 61% more and 14% less PEITC than the plants grown in 64 and 128 ppm sulfur, respectively. Increasing light intensity one week before harvest increased PEITC concentration of watercress leaves by approximately 60% in plants grown under a short day length (8 hours)

compared to those that were grown under continuous low light intensity.

Based on our research, we believe that growing watercress in media containing 200 ppm nitrogen and 128 ppm sulfur and providing supplemental lighting one week before harvest (during winter when the day lengths

"Studies are in progress to identify the environmental conditions and cultural methods that a greenhouse grower can adopt to enhance the nutraceutical content in these salad greens."

are short) may enhance the nutritional value of the harvested watercress crop.

In studies with purslane, we have found that plants grown with a nitrate (NO₃) to ammonium (NH₄) ratio of 0.5:0.5 contained about 241% and 53% higher alpha-linolenic acid compared to plants that were grown with a NO₃:NH₄ ratios 1:0 and 0.75:0.25 respectively. Providing the total nitrogen requirement in equal proportions of nitrate and ammonium forms appears to enhance the essential fatty acid content of purslane leaves.

Studies are in progress to identify the environmental conditions and cultural methods that a greenhouse grower can adopt to enhance the nutraceutical content in these salad greens and, thus, add value in terms of health benefits to the consumer.