

SUCCESS WITH HYDRANGEAS

Douglas A. Bailey, North Carolina State University

Greenhouse forcing of hydrangeas is not an easy task. However, a successful crop can be grown if a few key steps are followed. Listed below are some main points to keep in mind if you plan to grow hydrangeas.

Receiving and Establishing Pre-Cooled Plants

Hydrangeas are usually shipped in the late fall through early winter, after they have received a required cold storage treatment. They are received as dormant plants in 4-6" pots or as bare-root plants previously grown in 4" pots. Newly received plants should be allowed to initiate active root growth (for about 5 to 7 days) prior to transplanting into the final-sized pot. The ideal starting temperature for hydrangeas is a 60-62°F soil temperature supplied with bottom heat, while maintaining slightly cooler air temperatures (58°F). This allows root activity prior to buds opening on the shoots. Grow plants slightly on the "dry side" prior to transplant to prevent root rot and to encourage root development. No fertilizer should be applied until root activity and transplanting have occurred.

One of the main problems encountered with hydrangeas is poor root establishment leading to water stress damage during late stages of forcing. At transplant, the bottom of the root ball should be slit twice (to form an x pattern), about of the way up towards the top to form four sections; split open these sections and place them in direct contact with the soil in the new pot.

Flower Color Control


The key in assuring clear pink or blue inflorescences is ordering plant material programmed to develop the desired color and continuing the color program throughout forcing. Fertilization practices during the previous summer growth phase can affect coloration during forcing, and changing the color program during the forcing phase can result in interesting shades of mauve sometimes referred to as "blurple" tones.

Whether a hydrangea (excluding white cultivars) develops a pink or blue inflorescence is dependent on the presence and availability of aluminum. The absence of aluminum assures pink flowers; high availability of aluminum leads to blue flowers. By regulating aluminum, flower color can be controlled.

Cultivars vary in color tones and some are better suited for pinks while others are best produced as a blue. Select the cultivar with the color and tone best suited for your market demands. Although there are over 500 cultivars of hydrangeas, only a few are produced to any large degree (Table 1).

Pink Flowers. Avoid supplying aluminum to plants; do not use mineral soil in the substrate and use fertilizers that do not contain aluminum. Use relatively high levels of phosphorus in the fertilizer program. Phosphorus antagonizes aluminum uptake and helps assure pink flowers. Incorporate 3-4.5 lbs treble superphosphate (0-45-0) per yd³ into the substrate. Rotating mono-ammonium phosphate (11-53-00) into the feed program will also help raise phosphorus levels and help prevent aluminum uptake. An example feed program would be continuous feeding using 150 ppm nitrogen from 20-10-20 (10 oz/100 gal) rotated with 100 ppm nitrogen from 11-53-00 (18 oz/100 gal) every third feeding.

Try to maintain a substrate solution pH of 6.0-6.2; aluminum becomes more available at lower pH's. Be careful not to allow the pH to rise much above 6.4 or iron deficiency chlorosis will become a problem. If the pH of the irrigation water is higher than 6.5, consider acidifying to 6.3. Phosphoric acid would be the acidifier of choice for pink flowers, as it increases phosphorus levels in the substrate. If your irrigation water pH is low and the substrate pH declines with time, consider using calcium nitrate + potassium nitrate in the fertilizer rotation. Supply low to moderate levels of potassium. High levels of potassium tend to increase bluing of hydrangeas.



**Fine Perennial
Flowers,
Herbs, and
Native Plants
For The
Southern
Garden**

**P.O. Box 311
Lexington,
Georgia
30648**

GOODNESS GROWS
INCORPORATED

(706) 743-5055

The cultivars recommended for a dark pink to red are 'Böttstein' and 'Schenkenburg'; medium pinks include 'Merritt's Supreme', 'Kasteln', and 'Red Star'; and 'Rose Supreme' and 'Enziandom' produce light pink flowers. Some cultivars such as 'Mathilde Gütges' and 'Brestenburg' do not produce a consistent or clear pink and should be programmed as blue flowers only.

Blue Flowers. Although dormant plants purchased as blues will have received aluminum sulfate prior to shipment, aluminum must also be supplied during the forcing period. Start drenching with aluminum sulfate immediately after transplant. Apply 8 fl oz of drench per 6" pot using 10 lb aluminum sulfate/100 gallons of water. Drenches should be made to moist substrate only; drenching dry soil will result in damaged roots. Make applications at 10-14 day intervals. About 10 days after each application, measure the pH of the substrate. If the pH is higher than 5.6, another application of aluminum sulfate should be made. Continue this procedure throughout forcing. The aluminum sulfate not only supplies aluminum, it also maintains a low (5.2-5.5) pH in the substrate solution, desirable during forcing of blue hydrangeas. If the pH of the irrigation water is higher than 5.8, consider acidifying to 5.3. A 35% sulfuric acid source (should be available at auto supply stores) is the best acidifier for blue hydrangeas, as it will not add unwanted phosphorus and is not as caustic as a more concentrated acid.

Use a phosphorus-free substrate for transplanting and use a fertilizer lacking phosphorus. Apply high levels of potassium for increased bluing. For example, apply 150 ppm nitrogen and 300 ppm potassium at each irrigation supplied with ammonium nitrate (2oz/100 gal) plus potassium nitrate (11 oz/100 gal).

Height Control

Most hydrangeas, especially tall growing cultivars (Table 2), require height control during forcing. Apply B-Nine® sprays using 2500 ppm (most cultivars) to 5000 ppm (tall cultivars, especially 'Rose Supreme'). First applications are made when 3-5 leaf pairs have begun to unfold, about 3-4 weeks after the start of forcing. Under poor light conditions, repeat applications may be necessary at 10-14 day intervals. Treatments should be discontinued prior to when flower buds reach ¾" in diameter or inflorescences will be reduced in size at maturity. The following PGR treatments have been reported effective on hydrangeas, although they are not labeled: A-Rest® foliar sprays (50-100 ppm), A-Rest® substrate drenches (2-4 mg a.i./pot), Bonzi® sprays (50-100 ppm) Cycocel® drenches (0.3-1.6 g a.i./pot), and Sumagic® sprays (5-15 ppm).

Temperature and Timing

The rate of hydrangea development during forcing is directly related to average daily temperature, and to a certain degree, forcing speed can be regulated by adjusting temperature (Table 3). Generally, plants are forced in 80-100 days using 60°F nights/70°F cloudy day/75°F sunny day temperatures until flowers begin to show color (about 2½ weeks to sales date) then dropping the temperature until full coloration. At start of color, the temperature should be dropped to 54°F night/65°F day to intensify flower color.

Hardening and Post-Production Handling

At start of visible flower color, fertilization should be cut in half to help harden plants. Fully colored flowers are tender, and some shading to prevent overheating is beneficial during the last few weeks of production, especially for late crops such as for Mother's Day. Watering should be slowly reduced, but under no circumstances should plants be allowed to wilt.

Hydrangeas exhibit a long post-harvest life in the home if kept moist, out of direct light, and relatively cool. The key word for retailers and home owners is **Water**. Hydrangeas are very unforgiving when it comes to wilting and will never fully recover if allowed to dry out.

Pests and Diseases

Greenhouse pests and diseases differ from location to location. Listed below are the major problems of hydrangeas. Problems most likely to be encountered are indicated by the "helping hands":

Here's What's
HOT!

from
Flora Mart

<input type="checkbox"/> Adenium Obesum	<input type="checkbox"/> Mandevilla
<input type="checkbox"/> Dwarf Bougainvilleas	<input type="checkbox"/> Pentas
<input type="checkbox"/> Crotons	<input type="checkbox"/> Sword Fern
<input type="checkbox"/> Dwarf Cannas	<input type="checkbox"/> Scaevola "Blue Wonder"

Plus

★ HUNDREDS OF OTHER CUTTINGS & LINERS ★
For A Complete Price List Call Toll Free:
1-800-749-3148
"Your Best Source for
Plugs, Seeds, Cuttings & Liners"

5350 10th Ave. North • Suite 8
Lake Worth, Florida 33463
(407) 433-3157 / Fax: (407) 433-3952

Insects

Aphids (*Aphis gossypii*, *Myzus circumflexus*,
M. persicae)
Four-lined plant bug (*Poecilocapsis lineatus*)
Leaf-tiers (*Exartema ferriferanum*, *Udea*
rubigalis)
Rose-chaffer (*Macrodactylus subspinosus*)
Scale (*Lepidosaphes ulmi*, *Pulvinaria ssp.*)
Tarnished plant bug (*Lygus lineolaris*)
Thrips (*Hercinothrips femoralis*)
Whiteflies (*Bemisia tabaci*, *Trialeurodes*
vaporariorum)

Mites

Two-spotted mite or Red spider mite
(*Tetranychus urticae*)

Other Pests

Slugs (*Deroceras reficulatum*, *Limax spp.*)
Snails (*Helix spp.*)

Bacteria

Bacterial wilt (*Pseudomonas solanacearum*)

Fungi

Blister rust (*Pucciniastrum hydrangeae*)
Bud rot (*Botrytis cinerea*)
Gray mold (*Botrytis cinerea*)
Inflorescence blight (*Botrytis cinerea*)
Leaf spots (*Ascochyta hydrangeae*,
Cercospora arborescentis, *Corynespora*
cassicola, *Phyllosticta hydrangeae*, *Septoria*
hydrangeae)
Powdery mildew (*Erysiphe polygoni*)
Root rot (*Armillaria spp.*, *Polyporus spp.*,
Rhizoctonia spp., *Sclerotium spp.*)
Mycoplasma-Like Organisms (MLO)
Hydrangea virescence
Nematodes
Leaf nematodes (*Aphelenchoides spp.*)
Lesion nematodes (*Pratylenchus spp.*)
Root-knot nematodes (*Meloidogyne incognita*,
M. hapla)
Stem nematodes (*Ditylenchus dipsaci*)

Viruses

Alfalfa mosaic virus
Cucumber mosaic virus
Hydrangea mosaic virus
Hydrangea ring-spot virus
Tobacco rattle virus
Tobacco ring-spot virus
Tobacco necrosis virus
Tomato ring-spot virus
Tomato spotted-wilt virus

The Future For Hydrangeas

Forcing of hydrangeas for Valentine's Day is already common for many producers, and the trend seems to be increasing. The sales period for hydrangeas is definitely increasing and will continue to expand.

There is an increase in small (4½"-5" pots), single-stem hydrangeas. It should be possible to grow them and flower plants in a year-round program.


Another change which may take place in the future is an increase in the cultivars of the florists' hydrangea (*Hydrangea macrophylla* subsp. *macrophylla* var. *macrophylla*) and hydrangea species being produced. An example of "new" cultivars would be 'Kasteln'. A few years ago, it was relatively unknown; now it is becoming a major cultivar for late-season forcing.

With regards to new species, look for 'Pia' (*Hydrangea* x 'Pia') in the next few years. This selection is very dwarf, faster forcing than current cultivars, and very winter hardy. Pia is also noted because it appears to be a "non-bluing" cultivar that blooms pink regardless of how it is treated.

Other possible new hydrangeas include the lacecap varieties (*Hydrangea macrophylla* subsp. *macrophylla* var. *normalis*) such as 'Lybella', 'Blue Sky', 'Taube', 'Sunset', 'Shower', and 'Tricolor', which has variegated foliage. The lacecaps have a more open inflorescence and force more quickly than the florists' hydrangeas, you should consider experimenting with a few lacecaps this season.

KRUM

Horticultural Perlite



THE
IDEAL

SOIL
CONDITIONER

Krumbles the Earth

CAROLINA PERLITE COMPANY, INC.
Processors of Light Weight Aggregates Since 1949
P.O. BOX 158
GOLD HILL, NORTH CAROLINA 28071
(704) 279-2325

(Hydrangeas, continued)

Table 1. Description of the most common hydrangea cultivars.

Cultivar	Sepal color description		Relative days to flower*	Inflorescence size/plant height	Comments
	Grown as a pink	Grown as a blue			
Böttstein	Dark red	Not recommended	92	Medium/Short	
Brestenburg	Not recommended	Medium blue	83	Medium/Medium	Best medium blue
Enziandom	Light pink	Deep blue	92	Medium/Tall	
Kasteln	Dark pink	Medium blue	95	Medium/Short	Heat tolerant; good for late season forcing
Kuhnert	Not recommended	Medium blue	95	Small/Medium	
Mathilde Gütges	Not recommended	Light blue	95	Large/Medium	Best light blue
Merritt's Supreme	Dark pink	Medium blue	88	Large/Medium	Heat tolerant; good for late season forcing
Red Star	Light red	Medium blue	95	Large/Medium	
Rose Supreme	Light pink	Light blue	103	Very large/Very tall	Heat tolerant
Schenkenburg	Dark red	Not recommended	95	Medium/Medium	Best red
Sister Therese	White	White	95	Medium/Medium	Best white
Strafford	Dark pink	Not recommended	99	Large/Medium	Not heat tolerant; better for cool regions
Todi (Toddy)	Dark pink	Not recommended	88	Large/Medium	Not heat tolerant; better for cool regions

*Based on using a 60°F night/75°F day temperature up to 18 days prior to sale, then a 54°F night/65°F day for color intensification.

Table 2. Recommend rates of B-Nine based on cultivar height and responsiveness to B-Nine.

Cultivar	Relative height	Response to B-Nine	Recommended rate of B-Nine
Böttstein	Short	Great	0-2500 ppm
Enziandom	Tall	Moderate	5000 ppm
Kasteln	Short	Great	0-2500 ppm
Mathilde Gütges	Medium	Moderate	2500 ppm
Merritt's Supreme	Medium	Slight	2500-3500 ppm
Red Star	Medium	Very slight	5000 ppm
Rose Supreme	Very tall	Moderate	5000-7500 ppm
Schenkenburg	Medium	Moderate	2500 ppm

Table 3. Temperature effects on forcing timing. The table is based on cultivars that bloom in 88 days (such as 'Merritt's Supreme') when forced using a 60°F night/75°F day up to flower color. The timings given below assume that temperatures will be dropped to 54°F night/65°F day during the last 18 days of forcing.

Time interval	Night temperature/ day temperature (°F)		
	54/65	60/75	66/75
Days from start of forcing to bloom (salable color)	112	88	80
Days from start of forcing to pea-sized inflorescence (3/16" diameter bud)	43	32	28
Days from pea-sized bud to bloom	70	56	52
Days from nickel-sized inflorescence (13/16" diameter bud) to bloom	53	42	39
Days from silver dollar-sized inflorescence (1 1/2" diameter bud) to bloom	35	28	26
Days from first color (drop temperature to 54°F night/65°F day) to bloom	18	18	18



National Toll Free
1-800-749-3148

CURT SMITH

"Best Source for Plugs, Seeds, Cuttings & Liners"

Tel: (404) 299-1997

3519 Orchard Street
Clarkston, GA 30021